



**National
Oceanography Centre**
NATURAL ENVIRONMENT RESEARCH COUNCIL

UNIVERSITY OF
Southampton

Using R to Model Complex Biogeochemical Systems

Chris Wood

chris.wood@southampton.ac.uk

Introduction to the science

Introduction to the science

- Why are we interested in sediments?
 - 71% of Earth's surface (335,258,000 km²)


Introduction to the science

- Why are we interested in sediments?
 - 71% of Earth's surface (335,258,000 km²)
 - Very dynamic environments

Introduction to the science

- Why are we interested in sediments?
 - 71% of Earth's surface (335,258,000 km²)
 - Very dynamic environments
- Why are we interested in shelf-seas?

Introduction to the science




Introduction to the science

- Why are we interested in sediments?
 - 71% of Earth's surface (335,258,000 km²)
 - Very dynamic environments
- Why are we interested in shelf-seas?
 - Globally important sinks & sources for nutrients

Introduction to the science

- Why are we interested in sediments?
 - 71% of Earth's surface (335,258,000 km²)
 - Very dynamic environments
- Why are we interested in shelf-seas?
 - Globally important sinks & sources for nutrients
 - High rates of primary productivity



The maths...

The maths...

- Despite the complexity, it can be described mathematically(!):


The maths...

- Despite the complexity, it can be described mathematically(!):

$$\frac{d(1 - \emptyset)c_i}{dt} =$$

The maths...

- Despite the complexity, it can be described mathematically(!):


$$\frac{d(1 - \phi)c_i}{dt} =$$

Change in
concentration of i

The maths...

- Despite the complexity, it can be described mathematically(!):

$$\frac{d(1 - \emptyset)c_i}{dt} = \frac{\partial \left((1 - \emptyset)D_b \frac{\partial c_i}{\partial x} - \omega(1 - \emptyset)c_i \right)}{\partial x}$$

Change in
concentration of i

The maths...

- Despite the complexity, it can be described mathematically(!):

The diagram shows a mass balance equation for a chemical species i in a porous medium. The equation is:

$$\frac{d(1 - \phi)c_i}{dt} = \frac{\partial}{\partial x} \left((1 - \phi)D_b \frac{\partial c_i}{\partial x} - \omega(1 - \phi)c_i \right)$$

Annotations in the diagram include:

- A blue oval around the left-hand side term $\frac{d(1 - \phi)c_i}{dt}$, with an arrow pointing to the text "Change in concentration of i".
- A blue oval around the $(1 - \phi)D_b$ term inside the derivative on the right-hand side, with an arrow pointing to the text "Transport of chemical species".
- A blue oval around the $\omega(1 - \phi)c_i$ term inside the derivative on the right-hand side, with an arrow pointing to the text "Transport of chemical species".

The maths...

- Despite the complexity, it can be described mathematically(!):

$$\frac{d(1 - \phi)c_i}{dt} = \frac{\partial}{\partial x} \left((1 - \phi)D_b \frac{\partial c_i}{\partial x} - \omega(1 - \phi)c_i \right) - (1 - \phi)\Sigma(R(c_i, c_j))$$

Change in concentration of i

Transport of chemical species

The maths...

- Despite the complexity, it can be described mathematically(!):

The diagram shows a mass balance equation for chemical species i in a porous medium. The equation is:

$$\frac{d(1 - \phi)c_i}{dt} = \frac{\partial}{\partial x} \left((1 - \phi)D_b \frac{\partial c_i}{\partial x} - \omega(1 - \phi)c_i \right) - (1 - \phi)\sum_j R(c_i, c_j)$$

Annotations with arrows pointing to specific parts of the equation:

- Change in concentration of i** : Points to the left-hand side term $\frac{d(1 - \phi)c_i}{dt}$.
- Transport of chemical species**: Points to the divergence term $\frac{\partial}{\partial x} \left((1 - \phi)D_b \frac{\partial c_i}{\partial x} - \omega(1 - \phi)c_i \right)$.
- Rate of consumption of i , but dependent on j** : Points to the reaction term $(1 - \phi)\sum_j R(c_i, c_j)$.

The maths...

- Despite the complexity, it can be described mathematically(!):

$$\frac{d(1 - \emptyset)c_i}{dt} = \frac{\partial \left((1 - \emptyset)D_b \frac{\partial c_i}{\partial x} - \omega(1 - \emptyset)c_i \right)}{\partial x} - (1 - \emptyset)\Sigma \left(R(c_i, c_j) \right)$$

- Oxidic breakdown of organic matter:

The maths...

- Despite the complexity, it can be described mathematically(!):

$$\frac{d(1 - \emptyset)c_i}{dt} = \frac{\partial \left((1 - \emptyset)D_b \frac{\partial c_i}{\partial x} - \omega(1 - \emptyset)c_i \right)}{\partial x} - (1 - \emptyset)\Sigma \left(R(c_i, c_j) \right)$$

- Oxidic breakdown of organic matter:

$$R_{O_2} = \kappa[OM] \left(\frac{[O_2]}{[O_2] + \kappa S_{oxic}} \right)$$

Transport

Transport

- ReacTran package

Transport

- **ReacTran package**
- **written for ecological and biogeochemical models; the examples in the vignette reflect this**

Transport

- **ReacTran package**
- **written for ecological and biogeochemical models; the examples in the vignette reflect this**
- `Grid <- setup.grid.1D(N=100,dx.1=0.1,L=15)`

Transport

- **ReacTran package**
- **written for ecological and biogeochemical models; the examples in the vignette reflect this**
- `Grid <- setup.grid.1D(N=100,dx.1=0.1,L=15)`
 - `Grid$x.mid, Grid$x.int, Grid$dx`

Transport

- **ReacTran package**
- **written for ecological and biogeochemical models; the examples in the vignette reflect this**
- `Grid <- setup.grid.1D(N=100,dx.1=0.1,L=15)`
 - `Grid$x.mid, Grid$x.int, Grid$dx`
- `O2tran <- tran.1D (C=O2, C.up=bwO2, dx=Grid)`
 - **Other arguments allow specific transport terms to be used**

Solving the maths

Solving the maths

- The deSolve and rootSolve packages

Solving the maths

- The deSolve and rootSolve packages
 - rootSolve: ‘Nonlinear root finding, equilibrium and steady-state analysis of ordinary differential equations’

Solving the maths

- The deSolve and rootSolve packages
 - rootSolve: ‘Nonlinear root finding, equilibrium and steady-state analysis of ordinary differential equations’
 - deSolve: ‘General solvers for initial value problems of ordinary differential equations (ODE), partial differential equations (PDE), differential algebraic equations (DAE), and delay differential equations (DDE).’

Solving the maths (cont...)

Solving the maths (cont...)

- Simple to implement

Solving the maths (cont...)

- Simple to implement

```
modelFunction <- function(t, y, pars){  
  #implementation of transport and differential  
  #equations; e.g:  
  OC <- y[1:100]; O2 <- y[101:200]  
  oxicMin <- r*OC*(O2/(O2+ksO2oxic))  
}
```


Solving the maths (cont...)

– Simple to implement





```
modelFunction <- function(t, y, pars){  
  #implementation of transport and differential  
  #equations; e.g:  
  OC <- y[1:100]; O2 <- y[101:200]  
  oxicMin <- r*OC*(O2/(O2+ksO2oxic))  
}  
  
ss.output <- steady.1D(y=rep(10,2*100),  
  func=modelFunction, parms=c(r=10, ksO2oxic=1))
```

Solving the maths (cont...)

– Simple to implement

```
modelFunction <- function(t, y, pars){  
  #implementation of transport and differential  
  #equations; e.g:  
  OC <- y[1:100]; O2 <- y[101:200]  
  oxicMin <- r*OC*(O2/(O2+ksO2oxic))  
}  
  
ss.output <- steady.1D(y=rep(10,2*100),  
  func=modelFunction, parms=c(r=10, ksO2oxic=1))  
  
dyn.output <- ode.1D(y=ss.output$y, times=0:364,  
  func=modelFunction, parms=pars)
```

Output






Copyright © Ashley Hunn

Data management


- Data
 - Multiple sites, cruises, repeat measurements & parameters



ents

Data management

- Data
 - Multiple sites, cruises, repeat measurements & parameters
- R + MySQL (+ RJDBC/rJava)







Data management

- Data
 - Multiple sites, cruises, repeat measurements & parameters
- R + MySQL (+ RJDBC/rJava)
- Post-processing / data consistency checking carried out in R

Data management

- Data
 - Multiple sites, cruises, repeat measurements & parameters
- R + MySQL (+ RJDBC/rJava)
- Post-processing / data consistency checking carried out in R
- Allows model calibration to be carried out







Model testing & model calibration

Model testing & model calibration

- Sensitivity analysis
 - Allows us to discover the most sensitive model parameters

Model testing & model calibration

- Sensitivity analysis
 - Allows us to discover the most sensitive model parameters
- Genetic algorithm
 - (Relatively) efficient method of making model output fit real data



Future ideas / personal interests

Future ideas / personal interests

- Web-based model-runs via R + python
(mod_python / rpy2)

Future ideas / personal interests

- Web-based model-runs via R + python (mod_python / rpy2)
 - R would provide better graphics than currently available via PHP libraries

Future ideas / personal interests

- Web-based model-runs via R + python (mod_python / rpy2)
 - R would provide better graphics than currently available via PHP libraries
 - Greater access for non-modeller researchers

Future ideas / personal interests

- Web-based model-runs via R + python (mod_python / rpy2)
 - R would provide better graphics than currently available via PHP libraries
 - Greater access for non-modeller researchers
 - Public engagement of science

Any questions?

▪ Acknowledgements:

- Funding: **NERC & Defra**
- University of Southampton
 - Boris Kelly-Gerreyn, Peter Statham, Adrian Martin, Andy Yool, John Hemmings, Charlie Thompson, Carl Amos
- Netherlands Institute of Ecology
 - Karline Soetaert, Filip Meysman
- University of Portsmouth
 - Gary Fones, Fay Couceiro, Adam Hamilton
- Cefas
 - John Aldridge, Ruth Parker, Dave Sivyer, Johan van Molen, Naomi Greenwood, Elke Neubacher
- UEA
 - Keith Weston
- Partrac Ltd
 - Kevin Black, Rachel Helsby
- Crew of *Cefas Endeavour*